

Splenic Rupture

Ee Ling GOH*

Department of Emergency Medicine, Ng Teng Fong General Hospital, Singapore

Article Information

Received date: Apr 25, 2017

Accepted date: Apr 27, 2017

Published date: Apr 27, 2017

*Corresponding author

Ee Ling GOH, Department of Emergency Medicine, Ng Teng Fong General Hospital, 1 Jurong East Street 21, Singapore, 609606, Tel: +65 6716 2000; Fax: +65 6716 5500; Email: Ee_Ling_GOH@juronghealth.com.sg

Distributed under Creative Commons CC-BY 4.0

The Case

A 26 year old Indian man presented with recurrent left-sided abdominal pain and nausea to the emergency department. He had just returned from India few days ago and had just recovered from an upper respiratory tract infection. There was generalized abdominal tenderness. A bedside ultrasound revealed free fluid in the spleno-renal pouch. A Computed Tomography (CT) of abdomen and pelvis (Figure 1) showed presence of high-density fluid adjacent to the inferior pole of the spleen with haemoperitoneum. These findings were consistent with splenic rupture. The patient was subsequently tested positive for dengue.




Figure 1: High-density fluid adjacent to the inferior pole of the spleen.

Discussion

Dengue is a common mosquito-transmitted viral infection that is estimated to affect 50 to 100 million people worldwide each year [1]. Patients infected with dengue can have varied atypical and multisystemic presentations. To date, fewer than 15 cases of spontaneous splenic rupture due to dengue have been reported in the medical literature [2,3]. In these reported cases, the common clinical presentation was fever and abdominal pain with or without hemodynamic instability. Thrombocytopenia appeared to be a consistent feature. Splenectomy was performed for most of these patients. The pathogenesis of dengue haemorrhagic fever is unclear. The spleen is frequently congestive with subcapsular hematomas in 15% of cases [4]. This may predispose to spontaneous rupture in the absence of trauma. This can be misinterpreted as dengue shock syndrome which has a similar presentation. It is thus imperative to suspect and diagnose spontaneous splenic rupture which can be fatal. Management is dependent on the patient's hemodynamic status. Conservative treatment can be used but splenectomy is recommended for patients who deteriorate or needing more than 2 units of blood [3]. This case highlights the importance of recognizing this rare but potentially fatal complication of dengue by the emergency physician.

References

1. <http://www.who.int/mediacentre/factsheets/fs117/en/>
2. Seravali MR, Santos AH, Costa CE, Rangel DT, Valentim LF, Gonçalves RM. Spontaneous splenic rupture due to dengue fever: report of two cases. *Braz J Infect Dis*. 2008; 12: 538-540.
3. Rapp C, Debord T, Imbert P, Lambotte O, Roué R. Splenic rupture in infectious disease: splenectomy or conservative treatment? Report of three cases. *Rev Med Interne*. 2002; 23: 85-91.
4. Bhamarapravati N, Tuchinda P, Boonypaknavik V. Pathology of Thailandhemorrhagic fever. A study of 100 autopsy cases. *Ann Trop Med Parasitol*. 1967; 61: 500-510.